

AMENDMENTS TO THE CLAIMS:

Claims 7 and 14-15 are amended. The following is the status of the claims of the above-captioned application, as amended.

Claims 1-6 (Canceled).

Claim 7 (Currently amended). A method for reducing storage haze formation in a packaged tea extract, comprising:

- (a) contacting the tea extract with a pectin lyase;
- (b) separating insoluble solids from the tea extract; and
- (c) packaging the tea extract;

wherein the storage haze formation is reduced by at least 10% compared to an ~~untreated~~a tea extract not treated with a pectin lyase.

Claim 8 (Previously presented). The method of claim 7, wherein the storage haze formation is reduced by at least 50%.

Claim 9 (Previously presented). The method of claim 7, wherein the storage haze formation is reduced by at least 75%.

Claim 10 (Previously presented). The method of claim 7, wherein the storage haze formation is reduced by at least 90%.

Claim 11 (Previously presented). The method of claim 7, wherein the storage haze formation is reduced by at least 95%.

Claim 12 (Previously presented). The method of claim 7, wherein the storage haze formation is reduced by at least 99%.

Claim 13 (Previously presented). The method of claim 7, wherein the pectin lyase is a fungal pectin lyase.

Claim 14 (Currently amended). The method of claim 7, wherein the fungal pectin lyase is ~~derivable~~derived from *Aspergillus* sp.

Claim 15 (Currently amended). The method of claim 14, wherein the fungal pectin lyase is ~~derivable~~derived from *A.niger* or *A.oryzae*.

Claim 16 (Previously presented). The method of claim 7, wherein the amount of pectin lyase is in the range of from 0.1 to 1,000,000 UPTE per liter of the tea extract.

Claim 17 (Previously presented). The method of claim 16, wherein the amount of pectin lyase is in the range of from 1 to 100,000 UPTE per liter.

Claim 18 (Previously presented). The method of claim 17, wherein the amount of pectin lyase is in the range of from 10 to 10,000 UPTE per liter.

Claim 19 (Previously presented). The method of claim 18, wherein the amount of pectin lyase is in the range of from 1,000 to 8,000 UPTE per liter.

Claim 20 (Previously presented). The method of claim 7, wherein the pectin lyase is immobilized on a solid support.